

Microbial Genomics & Bioprocessing Research

Ching T. Hou
Microbiologist

Ph.D. Biochemistry, Tokyo University, Japan.
M.S. Biochemistry, Tokyo University, Japan.
B.S. Agriculture Chemistry, National Taiwan University, Taiwan

Microbial Genomics and Bioprocessing Research Unit
National Center for Agricultural Utilization Research
United States Department of Agriculture
Peoria, IL 61604-3999

Tel: (309) 681-6263
Fax: (309) 681-6672
E-mail: houct@ncaur.usda.gov

Research

Research in my laboratory focuses on developing new microbial processes for adding functionality groups to plant lipids or their component fatty acids to produce value-added food and industrial products. The goal is to develop new products from surplus soybean oil or its component fatty acids through bioprocesses. Oxygenated fatty acids such as hydroxy-, keto-, epoxy-, and cyclic fatty acids are important industrial chemicals and as materials in national defense. Castor oil and its derivatives (hydroxy fatty acid) are classified by the Department of Defence as strategic and critical materials. Newly discovered hydroxy unsaturated fatty acids were reported to have interesting physiological activity. Several microorganisms and biocatalytic processes have been identified that introduce new functional groups to the inner carbon atoms of fatty acids. These hydroxy fatty acids are classified into three types, namely: monohydroxy, dihydroxy and trihydroxy fatty acids.

DOD and TOD

A microbial culture, *Pseudomonas aeruginosa* PR3, was found to convert oleic acid to a new compound, 7,10-dihydroxy-8(*E*)-octadecenoic acid (DOD). The absolute configuration of DOD is 7(*S*),10(*S*)-dihydroxy-8(*E*)-octadecenoic acid. An intermediate for the synthesis of DOD was identified as 10*S*-hydroxy-8(*E*)-octadecenoic acid. DOD has surface-active properties and has antimicrobial activity against *Bacillus subtilis* and common clinical yeast, *Candida albicans*. Strain PR3 also converts ricinoleic acid to 7,10,12-trihydroxy-8(*E*)-octadecenoic acid (TOD). An intermediate, 10,12-dihydroxy-8(*E*)-octadecenoic acid was also identified for the synthesis of TOD. TOD showed anti-plant pathogenic fungal activities, especially on the rice blast fungus. Strain PR3 also converts linoleic acid to many isomeric 9,10,13 (9,12,13)-trihydroxy-11*E* (10*E*)-

octadecenoic acids. One kilogram of DOD was produced and several companies are investigating its application, uses include production of polyurethane rigid foam, skin care products and others.

12,13,17-THOA, THFAs, and DEOA

Another microbial culture, *Clavibacter* sp. ALA2 was found to convert linoleic acid to a new product, 12,13,17-trihydroxy-9(*Z*)-octadecenoic acid (12,13,17-THOA). 12,13,17-THOA inhibits the growth of many plant pathogenic fungi. Strain ALA2 also produced the following products from linoleic acid: two tetrahydrofuranyl fatty acids (THFAs), 12-hydroxy-13,16-epoxy-9(*Z*)-octadecenoic acid and 7,12-dihydroxy-13,16-epoxy-9(*Z*)-octadecenoic acid; two diepoxy bicyclic fatty acids, 12,17;13,17-diepoxy-9(*Z*)-octadecenoic acid (DEOA) and 12,17;13,17-diepoxy-7-hydroxy-9(*Z*)-octadecenoic acid (hDEOA). Tetrahydrofuranyl compounds are known to have anticancer activity. The diepoxy bicyclic fatty acids are complete new chemical entities. With many functionality groups on the molecules, their application in biomedical as well as specialty chemical industries is expected. We are trying to scale up production of these compounds. We are also working on lipases for production of structured lipids and healthy oils.

Selected Publications

Hosokawa, M., **Hou**, C.T., Weisleder, D., and Brown, W. 2003. Biosynthesis of tetrahydrofuranyl fatty acids from linoleic acid by *Clavibacter* sp. ALA2. *J. Am. Oil Chem. Soc.* 80:145–149.

Hou, C.T. 2003. New uses of vegetable oils: Novel oxygenated fatty acids by biotransformation. *SIM News.* 53:56–61.

Hosokawa, M., **Hou**, C.T., and Weisleder, D. 2003. Production of novel tetrahydrofuranyl fatty acids from α -linolenic acid by *Clavibacter* sp. Strain ALA2. *Appl. Environ. Microbiol.* 69:3868-3873.

Hosokawa, M., **Hou**, C.T., and Weisleder, D. Bioconversion of n-3 and n-6 polyunsaturated fatty acids by *Clavibacter* sp. ALA2. *J. Am. Oil Chem. Soc.* (in press, 2003).

Kim, H., Jang, Y.-S., and **Hou**, C.T. 2002. Effect of metal ions on the production of isomeric 9,10,13 (9,12,13)-Trihydroxy-11E(10E)-octadecenoic acid from linoleic acid by *Pseudomonas aeruginosa* PR3. *J. Enzyme Microbial Tech.* 30:752-757.

Iwasaki, Y., Brown, W., and **Hou**, C.T. 2002. Biosynthetic pathway of diepoxy bicyclic fatty acids from linoleic acid by *Clavibacter* sp. ALA2. *J. Am. Oil Chem. Soc.* 79:369-372.

Irimescu, R., Iwasaki, Y., and **Hou**, C.T. 2002. Study of TAG ethanolysis to 2-MG by immobilized *Candida antarctica* lipase and synthesis of symmetrically structured TAG. *J. Am. Oil Chem. Soc.* 79:879–883.

Lanser, A.C., Manthey, L.K., and **Hou**, C.T. 2002. Regioselectivity of new bacterial lipases determined by hydrolysis of triolein. *Curr. Microbiol.* 44:336-340.

Hou, C.T. 2002. Industrial uses of lipase. In *Lipid Biotechnology*. T.M. Kuo and Gardner, (eds) pp 387-397. Marcel Dekker, Inc. Press, New York.

Hou, C.T., Kuo, T.M., and Lanser, A.C. 2002. Value-added products through bioprocesses: New hydroxy fatty acids. *INFORM.* 13:307-316.

Kuo, T.M., Kim, H., and **Hou**, C.T. 2001. Production of a novel compound, 7,10,12-trihydroxy-8(E)-octadecenoic acid from ricinoleic acid by *Pseudomonas aeruginosa* PR3. *Curr. Microbiol.* 43:198-203.

Hou, C.T., Gardner, H.W., and Brown, W. 2001. 12,13,16-Trihydroxy-9(Z)-octadecenoic acid, a Possible intermediate in the biosynthesis of linoleic acid to tetrahydrofuran fatty acids by *Clavibacter sp.* ALA2. *J. Am. Oil Chem. Soc.* 78:1167-1169.

Kuo, T.M. and **Hou**, C.T. 2001. 7,10,12-Trihydroxy-8(E)-octadecenoic acid and derivatives and uses thereof. US Patent #6310007 B1.

Hou, C.T. 2000. Growth inhibition of plant pathogenic fungi by hydroxy fatty acids. *J. Ind. Microbiol. Biotechnol.* 24:275-276.

Kim, H., Gardner, H.W., and **Hou**, C.T. 2000. 10(S)-Hydroxy-8(E)-octadecenoic acid, an intermediate in the conversion of oleic acid to 7,10-dihydroxy-8(E)-octadecenoic acid. *J. Am. Oil Chem. Soc.* 77:95-99.

Hou, C.T. 2000. Lipases, Industrial Uses. *Encyclopedia of Microbiology.* 3:49-54.

Kim, H., Kuo, T.M., and **Hou**, C.T. 2000. Production of 10,12-dihydroxy-8(E)-octadecenoic acid, an intermediate in the conversion of ricinoleic acid to 7,10,12-trihydroxy-8(E)-octadecenoic acid by *Pseudomonas aeruginosa* PR3. *J. Ind. Microbiol. Biotechnol.* 24:167-172.

Hou, C.T. 2000. Biotransformation of unsaturated fatty acids to industrial products. *Advances in Appl. Microbiol.* 47:201-220.

Kim, H., Gardner, H.W., and **Hou**, C.T. 2000. Production of isomeric 9,10,13 (9,12,13)-trihydroxy-11E (10E)-octadecenoic acid from linoleic acid by *Pseudomonas aeruginosa* PR3. *J. Ind. Microbiol. Biotechnol.* 25:109-115.

Gardner, H.W., **Hou**, C.T., Weisleder, D., and Brown, W. 2000. Biotransformation of linoleic acid by *Clavibacter sp.* ALA2: Cyclic and bicyclic fatty acids. *Lipids.* 35:1055-1060.

Hou, C.T. 1999. Microbial production of a novel compound 7,10-dihydroxy-8-octadecenoic acid from oleic acid. US Patent # 5,900,496.

Gardner, H.W., and **Hou**, C.T. 1999. All (S) stereoconfiguration of 7,10-dihydroxy-8(E)octadecenoic acid from bioconversion of oleic acid by *Pseudomonas aeruginosa*. *J. Am. Oil Chem. Soc.* 76:1151-1156.

Hou, C.T., Gardner, H., and Brown, W. 1998. Production of multihydroxy fatty acids from linoleic acid by *Clavibacter sp.* ALA2. *J. Am. Oil Chem. Soc.* 75:1483-1487.

Hou, C.T. 1998. 12,13,17-Trihydroxy-9(Z)-octadecenoic acid and derivatives and microbial isolate for production of the acid. U.S. Patent # 5,852,196.

Hou, C.T., Brown, W., Labeda, D.P., Abbott, T.P., and Weisleder, D. 1997. Microbial production of a novel trihydroxy unsaturated fatty acid from linoleic acid. *J. Ind. Microbiol. Biotechnol.* 19:34-38.

Hou, C.T. 1997. Characterization of new yeast lipase. *J. Am. Oil Chem. Soc.* 74:1391-1394.

Hou, C.T. 1996. A novel compound, 12,13,17-trihydroxy-9(Z)-octadecenoic acid, from linoleic acid by a new microbial isolate *Clavibacter* sp. ALA2. *J. Am. Oil Chem. Soc.* 73:1359-1362.

Hou, C.T. 1995. A novel compound, 12,13,17-trihydroxy-9(Z)-octadecenoic acid from linoleic acid by a new microbial isolate *Clavibacter* sp. ALA2. *J. Am. Oil Chem. Soc.* 73:1359-1362.

Hou, C.T., Nakamura, L.K., Weisleder, D., Peterson, R.E., and Bagby, M.O. 1993. Identification of NRRL strain B-18602 (PR3) as *Pseudomonas aeruginosa* and effect of phenazine 1-carboxylic acid formation on 7,10-dihydroxy-8(E)-octadecenoic acid accumulation. *World J. Microbiol. Biotechnol.* 9:570-573.

Last Revised: April 6, 2005